

Jason K. Lowry, MD FAAOS

FL2145921

T0188705



ARLINGTON ORTHOPEDIC
ASSOCIATES, P.A.

LIFE IN FULL MOTION

800 Orthopedic Way

Arlington, TX 76015

p: (817) 375-5200

f: (817) 299-1789

2801 East Broad St

Mansfield, TX

76063 2001 N. MacArthur Blvd

Ste 630

Irving, TX 75061

<http://www.jasonlowrymd.com/>

<http://www.arlingtonortho.com/>

ACL RECONSTRUCTION (patellar tendon and hamstring graft) PROTOCOL

Last Modified: Oct 2012

<i>Preoperative</i>	<u>Goal</u>	<u>EXERCISE/METHODS</u>
Phase 1	Decrease swelling	CryoCuff with elevation (10-15 min)
	Restore terminal extension and flexion	Extension: prone hangs or heel props (7-10/min) Flexion: wall slides, heel slides, active assist flxn
	Normalize gait	Weight bearing as tolerated Treadmill: forward/backward (5-10 min)
	Normalize strength	Closed chain: leg press, total gym (10-50 deg), toe raises Open chain: Quad & hamstring machine (90-30 deg) 4-way SLR. All exercises 3-5 sets, 12-15 reps Electrical stimulation for VMO PRN
	Maintain aerobic fitness	Low-impact activities (bike, treadmill walking)
	Home program	Ice/elevation: 20 min on/60 min off Passive extension/flexion activities Lower extremity strengthening as indicated All activities 3-4 times/day or as indicated
Phase 2	Control swelling	Ace wrap with elevation-ankle pumps (20 min)
Postoperative Days 1-2	Restore terminal extension & flexion to 90	Extension: heel props (7-10 min) Flexion: heel slides (7-10 min) *avoid pillow under knee: lock-out brace in extension*
	Normalize gait	Weight bearing as tolerated

	Increase strength	SLR, quad sets, glute sets, hamstring sets
	Home program upon D/C	As above 4-5 times/day
Phase 3	Control swelling	Ace wrap with elevation-ankle pumps (20 min)
Postop	Restore terminal extension & flexion to 110 deg	Extension: prone hangs or heel props (7-10 min) Flexion: wall slides, heel slides, active assist flexion
Days 3-14	Normalize gait	Weight bearing as tolerated Treadmill: forward/backward (5-10 min) Calf, hamstring, hip flexor stretching (3 times, 30 sec hold) *emphasize heel to toe gait*
	Increase strength	4-way SLR, quad sets, multi-angle isometrics (0-45- 90 deg) Electrical muscle stimulation PRN (15 min) Partial squat (10-40 deg), toe raises with assistance PRN All exercises 3-5 sets, 12-15 reps Total Gym Level 1-3
	Home program	Ice/elevation: 20 min on/60 min off Passive extension/flexion activities Lower extremity strengthening as indicated All activities 3-4 times/day or as indicated *avoid pillow under knee: brace locked out in extension*
Phase 4	Control swelling	Ice PRN. Ace wrap or compression stockings PRN.
Postoperative		
Weeks 2-5	Full passive knee extension and flexion (0-125 by 4 wks)	Extension: prone hangs or heel props (7-10) Flexion: wall slides, heel slides, active assist flexion, bike (7-10 min)
	Normal ambulation	Treadmill: forward/backward (5-10 min) Increase grade with treadmill as tolerated Calf, hamstring, hip flexor stretching (3 times 30 sec hold)
	Pain-free strengthening	Closed chain: leg press (10-50), Total gym level (4-5) Push FROM without pain hamstring stretch, 4-way SLR, Toe raises, step ups, mini squats All exercises 3-5 sets, 12-15 reps

*Implement within painfree ROM once normal gait and effusion control are achieved
HS curls start at – 2 weeks for bone–patella–bone grafts and 4 weeks for HS grafts.

Initiate proprioception exercises (pain-free)

*BAPS (start with ball 1 sitting and progress to standing - 3-4 min)
*Double leg stand with rebounder
*Single leg stance/grid exercises (lunges, reaches) with eyes open/eyes closed
*Standing 4-way hip theratubing exercises
*Lateral treadmill walking both directions
*Implement once patient is able to ambulate normally without pain (3-5 min)

Criteria for progression to Phase 5:

**Phase 5
Postop
Weeks 6-12**

*No effusion
*Minimal crepitus.
*Full, Painless ROM
*Lysholm Score greater than 80 with no locking or instability

Improve aerobic level of fitness

Painfree low impact activity (bike, treadmill walking, stepper on own)
3-4 times per week, 20-30 minutes

Improve proprioception

Trampoline hopping bilaterally progressing To jogging, then single leg hopping (10 min)
Fitter with ski poles progressing to no ski poles (7-10 minutes) BAPS – single leg
Body blade Proprioceptive Star Exercises
*Week 10 – Light double leg hopping (multi-directional), lateral shuffles
*Only if all previous activities were completed without pain or increased swelling

Improve Strength

Total gym level 8-10 without pain, leg press (10-50), toe raises, step ups
Hamstring machine, 4-way SLR
All exercises 3-5 sets, 12-15 reps
Mini squats, Partial lunges, single-leg toe raise

Phase 6
Postop
Weeks 13-24

Criteria for progression to Phase 6:
*no effusion, painless full ROM,
*minimal crepitus.
*Lysholm score > 90 with no locking or instability
*Ratio of 80% involved to uninvolved on isokinetic test at 6 months post-op
*Ratio of 80% on one-legged jump

Improve aerobic level of fitness
(Usually 3-4 months)

Painfree low impact activity (bike, treadmill Walking, stepper on own) 3-4 times per Week, 20-30 minutes

Improve strength

Closed chain: leg press (10-50 deg), toe raises. Total gym level 8-10 without Pain.
Open chain: Quad machine (90-30 deg), Hamstring machine
All exercise's 3 sets 15-20 each

Implement running program

Walk/jog cycles
3-5 min warm-up (walk)
*10-30 sec jog/60 sec walk for 10-20 min 3-5 min cool-down (walk)
*increase jog time and decrease walk time based upon patient response.

Implement sport specific activity

*Progression program of strength training And agility/speed drills
*Program should be based upon specific patient needs.



Jason K. Lowry, MD