

Jason K. Lowry, MD FAAOS

FL2145921  
T0188705

[www.jasonlowrymd.com](http://www.jasonlowrymd.com)  
[www.arlingtonortho.com](http://www.arlingtonortho.com)



ARLINGTON ORTHOPEDIC  
ASSOCIATES, P.A.

LIFE IN FULL MOTION

Arlington Orthopedic Associates

800 Orthopedic Way  
Arlington, TX 76015  
p: (817) 375-5200  
f: (817) 299-1789

2801 East Broad St  
Mansfield, TX 76063

2001 N. MacArthur Blvd  
Ste 630  
Irving, TX 75061

## TOTAL HIP ARTHROPLASTY: DIRECT ANTERIOR APPROACH

PRIMARY & REVISIONS

### INPATIENT NURSING PROTOCOL

Last Revised: FEB 2015

1. 24 hours of IV ABX: Vanc + Ancef (or Clinda if PCN allergic)
2. DVT prophylaxis: early mobilization (RRP) is more important than chemoprophylaxis
  1. low-risk: ECASA 81mg BID x 30 days
  2. high-risk:
    1. normal renal fxn: Xarelto 10mg PO QDay x 30 days
    2. renal insufficiency: Lovenox 30mg SQ QDay x 30 days
3. ACTIVITY: UP OUT OF BED TO CHAIR WITHIN 2 HRS UPON ARRIVAL, THEN TID W/ MEALS
  1. "RRP" - Rapid Rehab Protocol
4. WB STATUS: GENERALLY WBAT w/ ROLLER WALKER
  - a. REVISIONS: MAY BE TOE-TOUCH DUE TO BONE LOSS \*\*THIS WILL BE CLEARLY NOTED IN THE ORDERS & OP REPORT\*\*
5. ANTERIOR HIP PRECAUTIONS X 4WKS
6. AN INTEGRAL PART OF "RRP" IS MULTI-MODAL PAIN MANAGEMENT WHICH INCLUDES:
  1. scheduled non-narcotic analgesics (eg acet, tramadol, & cox2s)
  2. minimal narcotic use
  3. regional anesthesia
  4. a local injection (cocktail, Exparel, etc)
  5. avoidance of PCA pumps.
7. Knee-high TED hose to bilateral LEs at all times x 4 wks
8. SCDs to bilateral LEs at all times while in the hospital
9. PILLOW: under operative knee while supine in bed; add b/t knees if side-lying
10. FOLEY: REMOVED IN PACU
  1. \*\*MALES >60y/o: FLOMAX PO QHS TO PREVENT URINARY RETENTION
11. DRAIN: CLAMPED X 2HRS POSTOP, DRAIN OUT ON POD1, PLACE STERILE 4X4 & TEGADERM, & THEN CHANGE PRN
12. Dressings:
  - a. DO NOT REMOVE AQUACEL: WATER-PROOF, SHOWERS OK, NO BATHS
    - i. CALL IF: > 60% STRIKE-THROUGH OR BROKEN SILICONE SEAL
    - ii. WILL BE REMOVED IN MY OFFICE AT POW2 VISIT (ALONG WITH STAPLES)
  - b. if NOT an aquacel, reinforce prn, then change on POD2 w/ same dressing
  - c. sutures/staples: as a general rule, there will NOT be staples or sutures to remove; a running subQ monofilament & derma bond will be used. \*\*

Dr. Jason K. Lowry, MD FAAOS