

phil h. berry, jr., m.d.
miguel a. hernandez, III, m.d.
ray f. aronowitz, m.d.
david a. heck, m.d.
zachary kelley, m.d.
alexander cho, m.d.
jason k. lowry, md
jon e. nathanson, d.p.m.



810 north zang blvd.
dallas, tx 75208-4233
telephone (214) 941-4243
fax (214) 941-1153

www.TheBoneDocs.com
www.jasonlowrymd.com

2800 east broad st, suite 510
mansfield, tx 76063
telephone (817) 453-3500
fax (817) 453-3820

ACL BRACE PROTOCOL (Post ACL surgery)

Last Modified: Oct 2012

Postoperative Brace

- Patients will be placed into a postoperative range of motion brace, locked in full extension.
- The range of motion brace may be unlocked for gait at 10-14 days (open 10 degrees greater than active ROM). The patient should be instructed on locking their brace out in full extension at night during sleep for the first 6 weeks to discourage flexion contracture.
- The patient will remain in this brace at all times, except when in therapy, or until the Don-Joy brace is given to them by orthopedics.

Functional Don-Joy Brace

- The patient will be measured for functional brace:
 - Preoperatively, if there is no significant swelling or quadriceps atrophy.
 - Postoperatively, at 8 weeks.
- Brace will be available in approximately 4 weeks.
- The functional brace may be worn no earlier than 6 weeks postoperatively.
- The patient must wear a brace at all times for the first 4 months after surgery.
- The brace may be discontinued for all activities of daily living, **but** must be worn for any activities involving sports with cutting, running on uneven surfaces.
- Athletes involved in contact sports should wear the functional brace indefinitely.

A handwritten signature in purple ink, appearing to read "J. K. Lowry".

Jason K. Lowry, MD