

phil h. berry, jr., m.d.  
miguel a. hernandez, III, m.d.  
ray f. aronowitz, m.d.  
david a. heck, m.d.  
zachary kelley, m.d.  
alexander cho, m.d.  
jason k. lowry, md  
jon e. nathanson, d.p.m.



810 north zang blvd.  
dallas, tx 75208-4233  
telephone (214) 941-4243  
fax (214) 941-1153

[www.TheBoneDocs.com](http://www.TheBoneDocs.com)  
[www.jasonlowrymd.com](http://www.jasonlowrymd.com)

2800 east broad st, suite 510  
mansfield, tx 76063  
telephone (817) 453-3500  
fax (817) 453-3520

## **Dr. Lowry's Post-Op Instructions**

### **Patient:**

1. Do not remove your splint/cast/dressings until I see you in clinic.
2. Keep your foot elevated at or, preferably, above heart level. This will dramatically help to decrease the expected swelling in your leg/ankle/foot. Remember: you can't elevate your foot too much & gravity is your enemy.
3. Apply two properly sealed ice packs to your foot / ankle / leg for 72 hours after your surgery. Large zip lock bags or small plastic trash bags work the best. Place one under & the other on top of the area. The key is to not get your splint/dressings wet. If you do, please call our clinic immediately.
4. Keep your splint/cast covered with a plastic trash bag during showers. Again, do whatever it takes to not get your splint/dressings wet.
5. Use your crutches at all times. It's OK to gently place your foot on the ground for balance. Otherwise, do not put any significant weight on your foot.
6. Please make an appointment to see me in 10-14 days after your surgery.

If you have any other questions or concerns please call my clinic.

Sincerely,

A handwritten signature in purple ink, appearing to read "J. Lowry", with a stylized flourish at the end.

Dr. Jason K. Lowry, MD FAAOS