

Jason K. Lowry, MD FAAOS

**FL2145921
T0188705**

www.jasonlowrymd.com
www.arlingtonortho.com



**ARLINGTON ORTHOPEDIC
ASSOCIATES, P.A.**

LIFE IN FULL MOTION

Arlington Orthopedic Associates

800 Orthopedic Way
Arlington, TX 76015
p: (817) 375-5200
f: (817) 299-1789

2801 East Broad St
Mansfield, TX 76063

2001 N. MacArthur Blvd
Ste 630
Irving, TX 75061

Dr. Lowry's Post-Op Instructions
LOWER EXTREMITY

DATE:

PATIENT:

DX:

PROCEDURE:

1. Do not remove your splint/cast/dressings until I see you in clinic.
2. Keep your foot elevated at or, preferably, above heart level. This will dramatically help to decrease the expected swelling in your leg/ankle/foot. Remember: you can't elevate your foot too much & gravity is your enemy.
3. Apply two properly sealed ice packs to your foot / ankle / leg for 72 hours after your surgery. Large zip lock bags or small plastic trash bags work the best. Place one under & the other on top of the area. The key is to not get your splint/dressings wet. If you do, please call our clinic immediately.
4. Keep your splint/cast covered with a plastic trash bag during showers. Again, do whatever it takes to not get your splint/dressings wet.
5. Use your crutches at all times. It's OK to gently place your foot on the ground for balance. Otherwise, do not put any significant weight on your foot.
6. Please make an appointment to see me in 14 days after your surgery.

If you have any other questions or concerns please call my clinic.

Sincerely,

Dr. Jason K. Lowry, MD FAAOS