

phil h. berry, jr., m.d.  
miguel a. hernandez, III, m.d.  
ray f. aronowitz, m.d.  
david a. heck, m.d.  
zachary kelley, m.d.  
alexander cho, m.d.  
jason k. lowry, md  
jon e. nathanson, d.p.m.



810 north zang blvd.  
dallas, tx 75208-4233  
telephone (214) 941-4243  
fax (214) 941-1153

www.TheBoneDocs.com  
www.jasonlowrymd.com

2800 east broad st, suite 510  
mansfield, tx 76063  
telephone (817) 453-3500  
fax (817) 453-3820

## **PLANTAR FASCIITIS REHAB PROTOCOL**

### **General**

- The plantar fascia originates off of the medial calcaneal tubercle, and the point of maximum pain is typically at that location.
- The treatment of plantar fasciitis is typically a protracted one, frequently lasting 6 months (or greater).

### **Phase I (Acute plantar fasciitis)**

- Shoe modifications - viscoheel cups, orthotics (soft or firm arch support).
- Activity modification - for runners – a period without running (usually from 2 to 6 weeks) until pain has subsided. When running resumed, begin with low mileage and increase slowly; run on soft surfaces – grass vs. pavement.
- Exercise
  - Ankle dorsiflexion stretches-towel stretches; wall stretches;block stretches (toes on block and heels on ground)
  - Extrinsic and intrinsic strength-towel scrunch; marble grab; toe raises
- Modalities
  - Iontophoresis
  - Phonophoresis
  - Fluidotherapy
  - Moist heat prior to activity/ Cryotherapy following activity

### **Phase II (Recalcitrant, if no improvement after 12 weeks of treatment)**

- Patient will be reevaluated by orthopaedist and possibly receive a steroid injection.
- If steroid injection is given; no running is permitted for at least 2 weeks.
- Patient resumes stretches, modalities.
- Night splinting is begun.

A handwritten signature in black ink, appearing to read "J. K. Lowry", with a stylized flourish at the end.

Jason K. Lowry, MD