



ARLINGTON ORTHOPEDIC
ASSOCIATES, P.A.

LIFE IN FULL MOTION

<http://www.jasonlowrymd.com/>

<http://www.arlingtonortho.com/>

RADIAL HEAD FRACTURE REHAB PROTOCOL

Introduction

- This protocol is recommended for Mason's type I fracture or type II or III with ORIF.
- **Goal:** Regain full pain-free ROM of elbow and prevent shoulder and wrist stiffness.

Phase (0 to 14 days)

- Begin elbow active ROM and active-assisted ROM for flexion and extension.
- ROM by the end of 2nd week should be at least 15 to 105 degrees.
- Begin putty/grip exercises.
- Begin isometric strengthening exercises for the elbow and wrist.

Phase II (15 days to 6 weeks)

- Continue elbow active and active assisted ROM exercises.
- Full flexion and extension ROM should be achieved by the end of 6 weeks.
- Begin active and active assisted supination and pronation.
- Begin light isotonic strengthening of flexion and extension.
- Assess and maintain shoulder and wrist strength and ROM.

Phase III (7-12 weeks)

- Continue active and active assisted supination and pronation.
- Full pronation and supination should be achieved by the end of the 8th week.
- Progressively increase isotonic strengthening in flexion/extension and pronation/supination.
- Work on any deficits.

Jason K. Lowry, MD